







SHORT COMMUNICATION article

Design of pyrrolidine-2,5-dione Schiff's base derivatives as GABA_A inhibitors for antiepileptic activity

Balsam A. Abodabous  , Suaad M. Abuskhuna  , Nisreen H. Meiqal  
Omran N.R. Fhaid *  

Department of Pharmaceutical and Medicinal Chemistry, Faculty of Pharmacy, University of Tripoli, Tripoli, Libya
* Author to whom correspondence should be addressed

Received: 29-03-2026, Accepted: 11-05-2026, Published online: 13-05-2026



Copyright © 2026. This open-access article is distributed under the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

HOW TO CITE THIS

Abodabous et al. Design of pyrrolidine-2,5-dione Schiff's base derivatives as GABA_A inhibitors for antiepileptic activity. *Mediterr J Pharm Pharm Sci.* 2026; 6(2): 36-43. [Article number: 251]. <https://doi.org/10.5281/zenodo.00000000>

Keywords: Anti-epileptic agents, GABA_A receptor, molecular docking, Schiff's base

Abstract: Succinimide derivatives are an essential class of compounds with diverse pharmacological applications. Different derivatives of succinimide as Schiff's base were designed for good inhibitory activity against the human GABA_A receptor. Compounds that inhibit GABA_A are effective as anti-epileptic agents. Virtual screening methods such as ADMET properties and Molecular docking were used. ADMET properties and drug likeliness of all investigated compounds were predicted by Swiss ADMET software. A higher binding affinity and interactions have been observed with the designed compounds compared to the standard compounds (Phenobarbital and Flumazenil).






Introduction


Substituted succinimides are important compounds of many drugs and drug candidates. One of the most fundamental objectives of organic and medicinal chemistry is the design and synthesis of molecules having value as human therapeutic agents. Cyclic imides and their derivatives contain an imide ring and the general structure of $-\text{CO}-\text{N}(\text{R})-\text{CO}-$, so they cross biological membranes *in vivo* [1]. There are several approved drugs with a cyclic imide structure. Substituted succinimide moiety appears as an interesting precursor of many biologically active such as CNS depressant [2], analgesic [3], antitumor and cytostatic [4, 5], anorectic [6], nerve conduction blocking antispasmodic [7], bacteriostatic [8], muscle relaxant and hypotensive [9], antibacterial [10], antifungal [11], anti-epileptic [12], and anti-tubercular [13, 14].

Although cyclic imide derivatives show a wide range of biological properties, the multidirectional activity of succinimide and its *N*-substituted derivatives prompted us to design several pyrrolidine-2, 5-dione Schiff's base derivatives (**Table 1**). Different methods have been reported for the synthesis of succinimide derivatives [15, 16] (**Scheme 1**). The designed compounds were used to study their binding interaction with the GABA-A receptor as antiepileptic agents by using the molecular docking technique (PyRx software). Evaluate their physicochemical properties, such as water solubility and lipophilicity, also predicting their pharmacokinetic parameters using the Swiss ADMET software.

General procedure for the synthesis of succinimide Schiff's base derivatives

Table 1: Succinimide Schiff's base derivatives

No	R	No	R
1		3	
2		4	
		6	

**Scheme 1:** Synthesis of pyrrolidine-2,5-dione Schiff's base derivatives

Materials and methods

Molecular docking: The docking studies of the designed (not synthesized) pyrrolidine-2, 5-dione Schiff's base derivatives have been carried out to predict the most possible type of interaction, the binding affinities, and the orientations of the docked ligands at the active site of the target proteins (GABA_A). The starting geometry of the pyrrolidine-2, 5-dione Schiff base derivatives was constructed using chem3D Ultra software (version 12.0, Cambridge Soft Com, USA). The optimized geometry of pyrrolidine-2, 5-dione Schiff's base derivatives with the lowest energy was used for molecular docking. Molecular docking was accomplished by Discovery Studio Visualizer overview 0.56. The crystal structures of Human GABA_A receptor alpha1-beta2-gamma2 subtype in complex with GABA and flumazenil (6D6T) and Human GABA_A receptor alpha1-beta2-gamma2 subtype in complex with GABA plus phenobarbital (6X3W) were downloaded from the Protein Data Bank (<http://www.rcsb.org/pdb>). The crystal structures of the GABA_A receptor (6D6T and 6X3W) were set up by cleaning the ligand and water molecules and saved as protein data bank (PDB) format through Discovery Studio (DS) software. Molecular docking was performed using PyRx software, with flumazenil and phenobarbital as GABA_A receptor control compounds. The polar hydrogen atoms were added to protein molecules. Then, partial atomic charges of the GABA_A receptor and designed compounds derivatives 1-6 molecules were calculated using Kollman methods. In the process of molecular docking, the grid maps of dimensions:(135.11Å X 121.58Å X 126.57 Å) and (94.06 Å X 112.79 Å X 110.56 Å) for 6D6T and 6X3 W, respectively. Discovery Studio software was used to analyze the docking results.

Results and discussion

Molecular docking analysis: Computer-aided drug designing, especially the molecular docking method, is an efficient technique to study the affinity and binding energy of a ligand with a studied receptor protein. The binding

affinity and energy of designed pyrrolidine-2,5-dione Schiff's base derivatives with the GABA_A receptor were obtained by using a molecular docking strategy [18-20].

According to the medical subject headings (MeSh) classification, phenobarbital and flumazenil are classified as GABA modulators, where they thought that GABA_A receptors have at least three allosteric sites at which these medications act [17, 21]. The docking energies of the reference compounds were utilized to compare the binding affinities of the tested compounds (**Table 2**). Three hydrogen bonds were observed between flumazenil with active sites by THR B: 122, LEU B: 128, and LYS B: 42. A single hydrogen bond was noticed between phenobarbital and the GABA_A receptor at the active site by THR B: 207. Other interactions of flumazenil and phenobarbital were found with the GABA_A receptor, such as π - π stacking and π - π -T-shaped interactions (**Figure 1**).




Figure 1: (i) Localization of flumazenil and phenobarbital (yellow shape) at active sites at GABA A
(ii) 2D interactions of drugs with amino acid residues

The molecular docking calculations tested compounds showed good binding affinity to the GABA_A receptor compared to the reference compounds (**Table 2**). Different bonds were observed in the binding of tested compounds **1 - 6**, the GABA_A receptor, such as H-bonding, Van der well forces, and electrostatic attraction. The observed interaction was between the aromatic ring of compounds **1 - 6** with Val 8: 292, ALA B: 328, LEU: 332, and LYS B: 71.

Compound **6** exhibited stronger interactions with active sites and with the highest binding affinities scores of the series GABA_A receptor (-9.1 and -8.0 Kcal/mol) for 6X3W and 6D6T, compared to standards (phenobarbital and flumazenil) with their binding affinity scores of (-6.3 and -6.5 kcal/mol), (-5.8 and -7.5 kcal/mol), respectively. The strength of binding is attributed to H-bonding with arginine amino acid 102 and more hydrophobic binding from the naphthyl group.

Compounds (**3** and **5**) with electron-donating groups (-OCH₃, -OH) show similar binding energies (-8.0 kcal/mol). The binding affinity of compound **2** to 6D6T (-7.7 kcal/mol) is higher than that of compounds **3** and **5**, with electron-donating groups. Introduction of a chlorine atom at the *p*-position on the aromatic ring has the same binding energy as compounds (**3** and **5**) with electron-donating groups.

Table 2: Binding energy of compounds 1 - 6 with GABA_A receptors










Compound	R	M. Wt	Docking energy (kcal/mol)	
			6x3 w	6D6T
Phenobarbital			-6.5	-6.3
Flumazenil			-7.3	-5.8
1		355.78	-8.1	-7.1
2		366.33	-8.1	-7.7
3		351.36	-8.0	-7.0
4		337.33	-8.0	-7.2
5		367.36	-7.7	-7.3
6		416.39	-9.1	-8.0

Table 3: (A) Distribution within the active site of the GABA_A receptor
(B) Predicted formed bonds with the GABA_A receptor

Comp. no	A	B
1		



Prediction of ADMET properties and drug-likeness

In-Silico ADMET prediction: To verify drug-likeness, canonical SMILES format of the designed compounds 1 - 6 were used in Swiss ADME (<http://www.swissadme.ch/index.php>) [22, 23]. The main bioavailability parameters were molecular weight (g/mol), lipophilicity (XlogP3), solubility (log S), polarity (Topological Polar

Surface Area - TPSA in Å²), saturation (fraction of carbon atoms in Sp³ hybridization-Csp³), and flexibility (No of rotatable bonds). Absorption parameters such as human intestinal absorption, blood-brain barrier, P-glycoprotein interaction (substrate or inhibitor), and metabolism (inhibitor or substrate) of the bio-active molecules with different cytochrome P450 enzymes, QSAR data for prediction of absorption, distribution, metabolism, excretion, and toxicity (ADMET) [24, 25].

Drug-like molecules, considering Ghose, Veber, Egan, Muegge, and Lipinski rules and good ADMET properties, were chosen as ligands in a subsequent molecular docking procedure. Molecular weight ≤ 500, hydrogen bond acceptor ≤ 10, hydrogen bond donor ≤ 5, Log P ≤ 5, Molar refractivity ≤ 140, comply with Lipinski's Rule of five and then log p o/w range between -2 to 6.5, polar surface area range between 7 to 200, log S (silicos-it) range lie above -4 and the drug score value above 0.5 is accepted one for designed compounds.

Table Error! No text of specified style in document.: ADME results of designed compounds

Comp	M.Wt	nHA	nRB	nHBD	nHBA	MR	TPSA (Å ²)	Log Po/w	Log S (SILICOS-IT)
1	355.78	25	5	1	4	97.66	78.84	2.52	-5.80
2	366.33	27	6	1	6	102.41	131.33	0.25	-4.56
3	351.36	26	6	1	5	99.14	88.07	1.95	-5.32
4	337.33	25	5	2	5	94.68	99.07	1.65	-4.62
5	367.36	27	6	2	6	101.17	108.30	1.69	-4.73
6	416.39	31	6	1	6	119.91	131.33	1.37	-6.20

Molecular weight: M.Wt, Number of heavy atom: nHA, Number of aromatic Heavy atom: nAHA, Number of sp³ hybridized carbon out of total carbon count: F. Csp³, Number of rotatable bonds: nRB, Number of H-bond acceptors: nHBA, Number of H-bond donors: nHBD, Molar refractivity: MR, topological polar surface area: TPSA

Pharmacokinetics and drug likeness: All investigated compounds showed high gastrointestinal absorption, indicating suitability for oral bioavailability. Regarding Cytochrome P450 (CYP450) inhibition, Compound 1 exhibited inhibitory effects on CYP1A2, CYP2C19, and CYP2C9 isomers, while showing no significant effect on CYP2D6 and CYP3A4. In comparison with compounds 2 - 6, compound 3 specifically inhibited CYP2C9.

Table 5: Pharmacokinetics parameters of the designed Schiff's base of Succinimide derivatives calculated with Swiss GI absorption

Comp	Absorption	BBB	P-gp Substrate	CYP1A2 inhibitor	CYP2C19 inhibitor	CYP2C9 inhibitor	CYP2D6 inhibitor	CYP3A4 Inhibitor	Log Kp (Skin Permeation) (cm/s)
1	high	no	no	yes	yes	yes	no	no	-7.14
2	high	no	yes	no	no	no	no	no	-8.06
3	high	no	no	no	no	yes	no	no	-7.69
4	high	no	no	no	no	no	no	no	-7.46
5	high	no	yes	no	no	no	no	no	-7.67
6	high	no	yes	no	no	no	no	no	-7.48

BBB permeant: P-gp: glycoprotein, CYP: Cytochrome P450 isomer

Drug-likeness prediction: Drug-likeness predictions were conducted based on established criteria, including Lipinski's Rule of Five parameters. According to Lipinski's Rule of Five, which suggests that absorption or permeation is more likely for molecules generally within accepted ranges, all designed compounds (1 - 6) met these criteria. Their molecular weights were typically under 500 g/mol. The Lipophilicity values (Log P o/w) for

the mono-substituted compounds ranged from 1.37 to 2.52. Furthermore, the compounds exhibited an optimum number of hydrogen bond donors (1 - 2) and acceptors (4 - 6) (Table 4).

The Ghose filter defines drug-likeness restrictions based on specific physicochemical parameters: a calculated log P value between -7.14 and -8.06, with the molecular weight (M. Wt) between 337.33 and 416.39, a molar refractivity between 97.66 and 119.91, and a total atom range of 25-31. All designed compounds 1-6 were found to comply with the established criteria for parameter assessment (Table 5).

The Veber rule subjects drug-likeness constraints as follows: a rotatable bond count ≤ 10 and a polar surface area (PSA) ≤ 140 . All the designed compounds (1 - 6) adhered to these parameters, further supporting their drug-like characteristics. All compounds (1 - 6) showed a similar bioavailability score of 0.55 (Table 5).

Table 5: Drug likeness of designed Schiff's base derivatives of succinimide calculated with Swiss ADME database

Comp	Lipinski	Ghose	Veber	Egan	Muegge	BA. Score
1	Yes	Yes	Yes	Yes	Yes	0.55
2	Yes	Yes	Yes	Yes	Yes	0.56
3	Yes	Yes	Yes	Yes	Yes	0.55
4	Yes	Yes	Yes	Yes	Yes	0.55
5	Yes	Yes	Yes	Yes	Yes	0.55
6	Yes	Yes	Yes	Yes	Yes	0.56

BA.S. Bioavailability score

Conclusion: A Series of pyrrolidine-2,5-dione Schiff base derivatives were designed against the GABA-A receptor. The compound exhibited high binding affinity toward the active site when compared to the reference compounds (flumazenil and phenobarbital). The nitro-substituted compounds (2 and 6) exhibited a strong binding affinity. Furthermore, the predicted ADMET properties indicated acceptable pharmacokinetic and toxicity properties. Based on the integration of molecular docking results, ADMET predictions, and reported biological activity, these pyrrolidine-2,5-dione Schiff base derivatives represent promising candidates for antiepileptic drug development. In particular, compounds 2 and 6 can be regarded as lead compounds for further structural optimization to enhance binding affinity and therapeutic efficacy toward the GABA-A receptor.

References

1. Alan Aitken R, Dheirya KS. The solid-state structures of cyclic NH carboximides. *Crystals*. 2020; 10(7): 606. doi: 10.3390/cryst10070606
2. Shivam AM, Deepika T, Panda D, Rahul R, Akhilesh KV. Visible light-promoted metal and oxidant-free stereo-selective synthesis of functionalized succinimides from aza-1,6-enynes. *RSC Sustainability*. 2025; 3: 592.
3. Saba Z, Muhammad H, Mirza WB. Synthesis, crystal structure, anti-cancer, anti-inflammatory, anti-oxidant, and quantum chemical studies of 4-(pyrrolidine-2,5-dione-1-yl) phenol. *Journal of Molecular Structure*. 2021; 1224: 129267. doi: 10.1016/j.molstruc. 2020.129267
4. Zefeng Z, Jiangxin Y, Xiaotong Ji, Meng N, Kaiwen K, Haifa Q, Xiaohui Z. Research progress in biological activities of succinimide derivatives. *Bioorganic Chemistry Maech*. 2021; 108: 104557.
5. Arun K, Anjali K, Prabhakar KV, Mano G, Girish Ch, Umesh K, Ashok KY, Deepak K. An insight into recent developments in imidazole-based heterocyclic compounds as anticancer agents: Synthesis, SARs, and mechanism of actions. *European Journal of Medicinal Chemistry*. 2024; 280: 15.116896.
6. Daniele RB, Laura VB, du VF, Rita DCV, Anelize D, Mariana Z, Thaise B, Priscila de RC. Antispasmodic activity of a series of cyclic imides (phthalimides) derivatives. *Naunyn-Schmiedebergs Arch Pharmacol*. 2026; 399(2): 2337-2352. doi: 1007/s00210-025-04515

7. Błażej G, Dariusz MP, Łukasz SZ. Succinimide derivatives as acetylcholinesterase inhibitors: In silico and in vitro Studies. *Curr Issues Mol. Biol.* 2024; 46(6): 5117-5130.
8. Sultan DYA, Dakhil ZM, Ali AA. Al. Antibacterial, antifungal, and antitumor properties of 2,5-pyrrolidinedione derivatives. *Chemistry Africa.* 2023; 6: 2933-2944. doi: 10.1007/s42250-023-00710-7
9. Muhammad IQ, Sami U, Umer R, Abdul S, Mater HM, Saif UKK, Muhammad MA. *N*-phenyl and *N*-benzyl substituted succinimide: Preclinical evaluation for their antihypertensive effect and underlying mechanism. *European Journal of Pharmacology.* 2024; 964(5): 176195.
10. Aeyaz AB, Nitin T, Iqbal S, Runjhun T. Structure-activity relationship (SAR) and antibacterial activity of pyrrolidine-based hybrids: A review. *Journal of Molecular Structure.* 2023; (1283)5: 135175.
11. Xin L, Yifang Ch, Yu W, Zhifu X, Ju P, Jixiang Ch. Design, synthesis, and antifungal activity of novel amide derivatives containing a pyrrolidine moiety as potential succinate dehydrogenase inhibitor. *Mol Divers;* 2024; 28(2): 805-816.
12. Krzysztof K, Sabina R, Jolanta O. Synthesis and anticonvulsant activity of new 1-[2-oxo-2-(4-phenylpiperazin-1-yl)ethyl]pyrrolidine-2,5-diones. *Bioorganic and Medicinal Chemistry Letters.* 2011; 21(19): 5800-5803.
13. Rakhi Y, Dilkhush M, Kavita S, Rajdeep T, Yogesh Y, Ram S. Recent advances in the synthesis of new benzothiazole-based anti-tubercular compounds. *RSC Adv.* 2023; 13: 21890-21925.
14. Braja GH, Vandana SP, Sanjeev KD, Suchitra D, Mahendra PD, Dharmendra S, Khanuja SPS. Bile acid amides derived from chiral amino alcohols: Novel antimicrobials and antifungals. *Bioorganic and Medicinal Chemistry Letters.* 2004; (14)3: 773-777.
15. Farshid H, Elham J. Cyclic imide derivatives: As promising scaffold for the synthesis of antimicrobial agents. *J Res Med Sci.* 2018; 23: 53. doi: 10.4103/ jrms. JRMS.539.17
16. Patil MM, Rajput SS. Succinimide: Synthesis, reaction and biological activity. *International Journal of Pharmacy and Pharmaceutical Sciences.* 2014; 6(11): 8-14.
17. Gallo AT, Hulse GK. Pharmacological uses of flumazenil in benzodiazepine use disorders: A systematic review of limited data. *Journal of Psychopharmacology.* 2021; 35(3): 211-220. doi: 10.1177/02698811209813
18. Alharati SH, Elbakay JAM, Hermann A, Gbaj AM. Polycystic ovary syndrome: Molecular modeling study on potential *Lepidium sativum* bioactive compounds in modulating kiss-1 gene function. *Mediterranean Journal of Medical Research.* 2025; 2(3): 129-140. doi: 10.5281/zenodo.17069661
19. Ieban NA, Gbaj HA, Gbaj MA, Hermann A, Gbaj AM. Immune profile score of modified Tremelimumab with CTLA-4 could enhance immunotherapy using molecular docking. *Mediterranean Journal of Medicine and Medical Sciences.* 2026; 2(1): 40-50. doi: 10.5281/zenodo.18667523
20. Hoque F, Nahar N, Annie FS, Rahim A, Hossain MK, Abdul Rahman MN, et al. Synthesis, characterization, and complexation of Schiff base ligand *p*-anisalcefuroxime with Cu²⁺ and Fe²⁺ ions; antimicrobial and docking analysis with PBP2xt to study pharmacokinetic parameters. *Mediterranean Journal of Pharmacy and Pharmaceutical Sciences.* 2025; 5(1): 48-64. doi: 10.5281/zenodo.14647651
21. Sherif FM. Pharmacological profile of the GABA-transaminase inhibitor vigabatrin. *World Journal of Pharmacy and Pharmaceutical Sciences.* 2015; 4(7): 139-148. doi: 10.20959/wjpps20157-4561
22. Daina A, Michielin O, Zoete V. Swiss ADME: A free web tool to evaluate pharmacokinetics, drug-likeness and medicinal chemistry friendliness of small molecules. *Sci Rep.* 2017; 7: 42717. doi: 10.1038/srep42717
23. Guan L, Yang H, Cai Y, Sun L, Di P, et al. ADMET-score - A comprehensive scoring function for evaluation of chemical drug likeness. *Med Chem Comm.* 2019; 10(1): 148-157.
24. Yang H, Lou C, Sun L, Li J, Cai Y, et al. ADMET SAR 2.0: Web-service for prediction and optimization of chemical ADMET properties. *Bioinformatics.* 2019; 35(6):1067-1069.
25. Tian S, Wang J, Li Y, Li D, Xu L, Hou T. The application of in silico drug-likeness predictions in pharmaceutical research. *Advanced Drug Delivery Reviews.* 2015; 86: 2-10. doi: 10.1016/j.addr.2015.01.009

Author's contribution: ONRF conceived and designed the study. SMA collected data. ONRF & NHM contributed to data analysis, and BAA performed the data analysis and interpretation. SMA drafted the manuscript. All authors approved the final version of the manuscript and agreed to be accountable for its contents.

Conflict of interest: The authors declare the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Ethical issues: The authors have considered ethical issues, including plagiarism, informed consent, data fabrication or falsification, and double publication or submission.

Data availability statement: The raw data that support the findings of this article are available from the corresponding author upon reasonable request.

Generative AI disclosure: No Generative AI was used in the preparation of this manuscript.